

WHAT IS CLAIMED IS:


- 5 1. A method of promoting the rate of hematopoietic cell multiplication, comprising administering an effective amount of a CXCR4 antagonist to hematopoietic cells.
- 10 2. The method of claim 1, wherein the hematopoietic cells are hematopoietic stem or progenitor cells.
- 15 3. A method of increasing the circulation of hematopoietic cells in a patient in need of such treatment, comprising administering to the patient an effective amount of a CXCR4 antagonist to mobilize the hematopoietic cells from a marrow locus to a peripheral blood locus.
- 20 4. The method of claim 1, further comprising introducing a heterologous gene into the hematopoietic cells for gene therapy.
5. The method of claim 1, wherein the hematopoietic cells are *ex vivo*.
- 25 6. The method of claim 1, wherein the hematopoietic cells are *in vivo*.
7. The method of claim 1, wherein the hematopoietic cells are selected from the group consisting of hematopoietic stem cells and hematopoietic progenitor cells (including CFU-GEMM, BFU-E, CFU-Meg, CFU-GM, CFU-M/DC CFU-E₀, CFU-Bas, Pro-B cells and lymphoid stem cells), that are known to differentiate into mature myeloid and lymphoid blood cells, including erythrocytes, platelets, neutrophils, monocytes, macrophages, dendritic cells (myeloid and lymphoid related), eosinophils, basophils, mast cells, B cells. and T cells.
- 30 8. The method of claim 1, wherein the CXCR4 antagonist comprises a CXCR4 antagonist peptide.

9. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

5	KGVSLSYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQ VCIDPKLKWIQEYLEKALN (SEQ ID No. 1);
	KGVSPSYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQ VCIDPKLKWIQEYLEKALN (SEQ ID No. 2);
10	KGVSLPYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQ VCIDPKLKWIQEYLEKALN (SEQ ID No. 3);
	KGVSLSPRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQ VCIDPKLKWIQEYLEKALN (SEQ ID No. 4);
15	KGVSLSYPCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQ VCIDPKLKWIQEYLEKALN (SEQ ID No. 5);
	KGVSP*SYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 6);
20	KGVSLP*YRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 7);
	KGVSLSP*RCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 8);
	KGVSLSY*PCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 9);
30	KGVSBtdYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 10);
	KGVSLBtdRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 11);
35	KGVSLSBtdCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNR QVCIDPKLKWIQEYLEKALN (SEQ ID No. 12);


40

wherein P* =



with X= Ar, Ar-OH, alkyl and more

and Btd =



X= Alkyl, Ar, Ar-OH and more

5

10. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

- a) KGVLSYRCPCRFFESH
- b) KGVLSYRC

10

11. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

- KGVS**P**SYRCPCRFFESH (SEQ ID No. 17)
- KGVS**L**PYRCPCRFFESH (SEQ ID No. 18)
- KGVS**L**SPRCPCRFFESH (SEQ ID No. 19)
- KGVS**L**SY**P**CPCRFFESH (SEQ ID No. 20)
- KGVS**P***SYRCPCRFFESH (SEQ ID No. 21)
- KGVS**L****P***YRCPCRFFESH (SEQ ID No. 22)
- KGVS**L****S****P***RCPCRFFESH (SEQ ID No. 23)
- KGVS**L**SY**P***CPCRFFESH (SEQ ID No. 24)
- KGVS**Btd**YRCPCRFFESH (SEQ ID No. 25)
- KGVS**L****Btd**RCPCRFFESH (SEQ ID No. 26)
- KGVS**L**SBtdCPCRFFESH (SEQ ID No. 27)
- KGVS**P**SYRC (SEQ ID No. 28)
- KGVS**L**PYRC (SEQ ID No. 29)
- KGVS**L**SPRC (SEQ ID No. 30)
- KGVS**L**SY**P**C (SEQ ID No. 31)
- KGVS**P***SYRC (SEQ ID No. 32)
- KGVS**L****P***YRC (SEQ ID No. 33)

15


20

25

KGVSLSP*RC (SEQ ID No. 34)
 KGVSLSYPC (SEQ ID No. 35)
 KGVSBtdYRC (SEQ ID No. 36)
 KGVSLBtdRC (SEQ ID No. 37)
 KGVSLSBtdC (SEQ ID No. 38)

5


wherein P* =



with X= Ar, Ar-OH, alkyl and more

10

and Btd =



X= Alkyl, Ar, Ar-OH and more


15

12. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSPSYRC	KGVSLPYRC	KGVSLSPRC	KGVSLSYPC
KGVS P SYRC	KGVS L PYRC	KGVS L SPRC	KGVS L SYPC
KGVSP*SYRC	KGVSLP*YRC	KGVSLSP*RC	KGVSLSYPC
KGVS P *SYRC	KGVSL P *YRC	KGVSLSP*RC	KGVSLSYPC
KGVS Btd YRC	KGVSL Btd RC	KGVSLSBtdC	
KGVS Btd YRC	KGVSL Btd RC	KGVSLSBtdC	

20


wherein P* =



with X= Ar, Ar-OH, alkyl and more

5


and Btd =



X= Alkyl, Ar, Ar-OH and more

10


13. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:



wherein X is a natural or unnatural amino acid linker between each of the arginines at position 8 in each sequence; and,

15


wherein P* =



with X= Ar, Ar-OH, alkyl and more

5

and Btd =



X= Alkyl, Ar, Ar-OH and more

10

14. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVLSYRCP CRFF-G_n-LKWIQEYLEKALN (SEQ No. 63)

KGVLSYRCP CRFFESH-G_n-LKWIQEYLEKALN (SEQ No. 64)

15

wherein n is an integer from 0 to 10.

15. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVLSYRCP CRFF-(CH₂)_n-LKWIQEYLEKALN (SEQ No. 65)

20

KGVLSYRCP CRFFESH-(CH₂)_n-LKWIQEYLEKALN (SEQ No. 66)

where n is an integer from 1 to 20.

16. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

25

KGVSPSYRCP CRFF-GGGG-LKWIQEYLEKALN;

KGVSLPYRCP CRFF-GGGG-LKWIQEYLEKALN;

KGVSLSPRCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSLSYPCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSPSYRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLPYRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 5 KGVSLSPRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLSYPCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSPSYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLPYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSPRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 10 KGVSLSYPCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSPSYRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLPYRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSPRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSYPCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN,

15

wherein n is an integer from 1 to 20.

17. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

20 KGVSL¹*SYRCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL²*YRCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL³*RCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSLSY⁴*CPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL⁵*SYRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 25 KGVSL⁶*YRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLSP⁷*RCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSLSY⁸*CPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSL⁹*SYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL¹⁰*YRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 30 KGVSLSP¹¹*RCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSY¹²*CPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL¹³*SYRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL¹⁴*YRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;

KGVSLSP^{*}RCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSLSY^{*}RCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;

5

KGVSL^{Btd}SYRCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL^{Btd}RCPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL^{Btd}CPCRFF-GGGG-LKWIQEYLEKALN;
 KGVSL^{Btd}SYRCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSL^{Btd}RCPCRFFESH-GGGG-LKWIQEYLEKALN;
 KGVSL^{Btd}CPCRFFESH-GGGG-LKWIQEYLEKALN;


10

KGVSL^{Btd}SYRCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL^{Btd}RCPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL^{Btd}CPCRFF-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL^{Btd}SYRCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL^{Btd}RCPCRFFESH-(CH₂)_n-LKWIQEYLEKALN;
 KGVSL^{Btd}CPCRFFESH-(CH₂)_n-LKWIQEYLEKALN,

15

wherein n is an integer from 0 to 20 and


wherein P^{*} =



with X = Ar, Ar-OH, alkyl and more

20

and Btd =



X = Alkyl, Ar, Ar-OH and more

25

18. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCP~~CRFFGGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFESHGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFGGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFESHGGG~~GLKWIQEYLEKALN



- 5 19. A CXCR4 antagonist peptide selected from the group consisting of:

KGVSLSYRCP~~CRFFGGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFESHGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFGGGG~~GLKWIQEYLEKALN



KGVSLSYRCP~~CRFFESHGGG~~GLKWIQEYLEKALN



- 10 20. The method of claim 8, wherein the CXCR4 antagonist peptide is selected from the group consisting of:

KGVSLSYRCP~~CRFFGGGG~~SKPGVIFLTKRSRQV;

KGVSLSYRCP~~CRFF(CH₂)_n~~SKPGVIFLTKRSRQV;

KGVSLSYRCP~~CRFFGGGG~~EEWVQKYVDDLELSA;

KGVSLSYRCP $\text{CRFF}(\text{CH}_2)_n$ EEWVKYVDDLELSA,

where n is 0 or an integer between 1 and 20.

- 5 21. A method of treating a cancer in a patient in need of such treatment comprising administering an effective amount of a CXCR4 antagonist to the patient to promote the rate of hematopoietic cell multiplication.
- 10 22. A method of treating an autoimmune disease in a patient in need of such treatment comprising administering an effective amount of a CXCR4 antagonist to the patient to promote the rate of hematopoietic cell multiplication.

15